



CORONA MASTER™

8-27-0

Guaranteed analysis

Total nitrogen (N): 8%
6% Ammoniacal nitrogen (N)
2% Urea nitrogen (N)
Available Phosphate (P₂O₅): 27%
Magnesium (Mg): 0.9 %
Sulfur (S): >7,2%
Boron (B)..... 0.5%
0.5% Water Soluble Boron (B)
Copper (Cu) 1%
1% Water Soluble Copper (Cu)

Manganese (Mn)..... 6%
6% Water Soluble Manganese (Mn)
Molybdenum (Mo)..... 0.1%
0.1% Water Soluble Molybdenum (Mo)
Zinc (Zn)..... 5%
5% Water Soluble Zinc (Zn)
DERIVED FROM Monoammonium phosphate,
Manganese sulfate, Boron pentahydrate, Zinc
sulfate, Magnesium sulfate, Urea, Ammonium
sulfate, Copper sulfate, Sodium molybdate.

F1550

PLEASE READ CAREFULLY BEFORE USING

KEEP OUT OF REACH OF CHILDREN

Crop	Nº of applications	Application stages	Application rate
All row crops including Soybeans, Corn, Wheat, Grains, Peanuts, all potatoes	1 – 3	Developmental and post bloom	2 – 4 lb/A
Citrus	1 – 10	Pre bloom , post bloom, summer, fall spray	1 – 10 lb/A
All fruits and nut crops	1 – 10	Pre bloom, bloom, summer and fall flushes	1 – 10 lb/A
All vegetables	1 – 10	Development stages ,pre bloom, correct deficiencies	1 – 10 lb/A
Turfgrasses and Ornamentals	1 – 3	For plant establishment	2 – 3 lb/A

INDICATIONS DOSAGE AND CONDITIONS OF USE: Add CORONA Master first in a half tank water and agitate while mixing. Spray Oils should be added very last to any mix so water solubles can dissolve completely first. Keep agitating until the end of the application. Dilute into at least 15-20 gallons/Acre. Foliar fertilization is intended as a supplement to a regular fertilization program and may not by itself provide all the nutrients normally required by crops or other intended plants. Do not spray during periods of frost. In hot conditions spray at the beginning or at the end of the day.

COMPATIBILITY AND STORAGE: CORONA Master is compatible with many chemicals. In all cases, test the compatibility prior to treatment. Storage: Keep out of moisture.

Warning: Some crops may be injured by application of Boron. This product contains molybdenum (Mo). Crops high in molybdenum can be toxic to ruminant animals. Use only according to manufacturer's directions.

Net Weight: 13.22 lbs (6 kg)

Timac Agro USA, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Timac Agro USA, Inc.. In no case shall Timac Agro USA, Inc. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the Buyer. Timac Agro USA, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

Timac Agro USA, INC. – P.O. Box 888 - Route 724 & I-176 Reading, PA 19607-0888 - USA

DANGER - H318: Causes serious eye damage. H360: May damage fertility or the unborn child. H373 - May cause damage to organs (brain) through prolonged or repeated exposure (Inhalation). P201: Obtain special instructions before use. P260: Do not breathe dust. P280: Wear face shield, protective clothing, protective gloves. P305+P351+P338: If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. P308+P313: If exposed or concerned: Get medical advice/attention. P310: Immediately call a doctor, a POISON CENTER.

Manufactured in Spain by: Timac Agro España
S.A. Polígono Arazuri-Orcoyen - Calle C, nº32
ORCOYEN (NAVARRA)
31160 ESPAÑA - Tel: + 34 948 324 500

2EACOROMA-USA6 2039 - Made in Spain by Timac Agro España