EXCELIS MAXX RHIZOVIT



CONTAINS NON-PLANT FOOD INGREDIENTS

Active Ingredients:

86% Other Ingredients

F1550

EXCELIS MAXX® fertilizer additive is designed to protect urea-based fertilizers from ammonia volatilization with no crop restrictions. This product is recommended for surface or shallow-placed (less than two inches) urea and urea-containing fertilizers. It can be used pre-plant, pre-emergence, side-dress, top-dress or other post-planting applications. This product is not plant food and is not intended for use as a fertilizer.

COMPATIBILITY: EXCELIS MAXX is compatible with most agricultural fertilizers and blends, however jar tests are recommended prior to the preparation of mixtures. It is also recommended that **EXCELIS MAXX**/UAN spray mixtures with pesticides, adjuvants, and (or) other products be applied to the soil as soon as possible after preparation.

STORAGE AND HANDLING: EXCELIS MAXX should be stored in a temperature range of 45°-100°F. Store in original container only. Keep packages tightly closed and do not allow water to be introduced to the container. Do not store near heat or an open flame. Exposure of **EXCELIS MAXX** to below freezing temperatures for an extended period may result in the formation of gel and/or crystals. If this occurs, expose the product to ambient temperatures greater than 55°F. This will result in reconstitution of the product, with no degradation of the active ingredient. Always wear appropriate chemical mixing personal protective equipment (PPE), including eye/face and hand protection.

DENSITY: 10.0 lbs/gal at 68°F

USE RATES:

- 32 Fl Oz Per Ton Of Granular Urea
- 25 Fl Oz Per Ton Of Liquid UAN

PLEASE READ CAREFULLY BEFORE USING - KEEP OUT OF THE REACH OF CHILDREN.
RECIRCULATE WELL BEFORE USE. STORE ABOVE 45°F

DANGER

H315: Causes skin irritation.

H317: May cause an allergic skin reaction

H318: Causes serious eve damage

H361: Suspected of damaging fertility (if swallowed, in contact with skin, if inhaled).

P202: Do not handle until all safety precautions have been read and understood.

P280: Wear face protection, protective clothing, protective gloves.

P302+P352: If on skin: Wash with plenty of soap and water.

P305+P351+P338: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

P310: Immediately call a doctor or POISON CENTER.

P308+P313: If exposed or concerned: Get medical advice/attention.

DIRECTIONS FOR USE:

Several environmental factors contribute to the degree and rate of nitrogen loss due to volatility, denitrification, and leaching. When considering the appropriate rate of EXCELIS MAXX, factors known to increase loss should be considered:

- 1) Very wet soils.
- **2)** The intensity of sunlight, wind, and humidity.
- 3) Conditions of high heat (both air and soil).
- **4)** Higher soil pH levels have a greater potential for loss due to volatility.
- **5)** Soil organic matter (OM), cation exchange capacity (CEC), and sandy soils (low levels of OM or CEC) increase nitrogen loss potential.
- 6) High amounts of crop residue or stubble tend to increase loss due to higher urease layels

When conditions for increased nitrogen loss exist, or a longer length of nitrogen stabilization is needed, the rate may be increased by 25%-50%.

PREPARATION OF EXCELIS MAXX + UREA BLENDS:

For direct blending/impregnation on urea, a fertilizer blender or equipment capable of mixing urea directly with the product is required. Rotary, auger, and paddle mixers are all applicable – but their surfaces must be clean, dry, and free of rust. Weigh the urea and transfer it directly to the blender. Measure the desired amount of EXCELIS MAXX and apply directly to the urea. Blend the urea/EXCELIS MAXX mixture until coverage is uniform, EXCELIS MAXX contains a blue dye to help in the verification that blending is sufficient. Under certain temperature/humidity conditions, **EXCELIS MAXX**-treated fertilizer may clump together in the blender due to moisture. This situation may be prevented by including other types of fertilizers (e.g., monoammonium phosphate/diammonium phosphate, potash, etc.) or by adding drying agents based on attapulgite clay. Include the drying agent at the rate of 0.25-0.5% of the urea/EXCELIS MAXX.

PREPARATION OF EXCELIS MAXX + UREA-AMMONIUM-NITRATE (UAN) BLENDS:

Fill the spray application tank to half of its volume with UAN. While maintaining agitation, add the desired amount of **EXCELIS MAXX** and mix well. Complete the mixture by adding the remainder of the UAN.

CONDITIONS OF SALE-LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES:

If these terms are not acceptable, return the product, unopened, and the full purchase price will be refunded. This label is believed to be reliable and must be followed carefully. Injury to crops, animals, humans, or the environment where the product has been applied may result from extraordinary or unusual weather conditions, failure to follow the label directions, or poor application practices, all of which are beyond the control of TIMAC AGRO USA ("the manufacturer") and the seller. The manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to. The manufacturer makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any purpose, and no such warranty shall be implied by law. The exclusive remedy against the manufacturer or seller for any cause of action relating to the handling or use of this product shall be limited to, at the seller's choice, one of the following: 1. Refund of the purchase price, or 2. Replacement of the product used. Neither the manufacturer nor the seller shall be limited to, or on special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income by the end user. The manufacturer and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

Manufactured By: TIMAC AGRO USA INC P.O. Box 888 - Route 724 & I-176 Reading, PA 19607-0888 - USA



This product is not plant food and is not intended for use as a fertilizer.



