



FERTILEADER Elite - 9-0-6

GUARANTEED ANALYSIS

• Total Nitrogen (N)...... 9 % Nitrate Nitrogen • Soluble Potash (K₂O)6% • Calcium (Ca)8.5%

DERIVED FROM: Calcium nitrate, Potassium hydroxide, Potassium nitrate, Sodium octaborate,

F 1550

NON PLANT FOOD INGREDIENTS:

Plant and seaweed extracts...... 4 %

DIRECTIONS FOR USE:

PLEASE READ CAREFULLY BEFORE USING - KEEP OUT OF REACH OF CHILDREN - RECIRCULATE BEFORE USE

The use of FERTILEADER Elite helps to increase sugars. The general time for this activity is 14 to 36 days after application. Do not spray during periods of frost. In hot conditions it is better to spray at the beginning or at the end of the day. Add FERTILEADER Elite in a half clean sprayer and agitate while mixing. Keep agitating until the end of the application. Dilute into at least 20 gallons/Acre. It can also be used in ULV aerial application. Foliar fertilization is intended as a supplement to a regular fertilization program and may not by itself provide all the nutrients normally required by crops or other intended plants.

Crop	N° of applications	Application stages	Application rate pts/Acre
Melon, Watermelon	1 – 2	40 to 50 days after planting and 10 days later	2-2.5
Tomato, Pepper, Eggplant	3 – 5	From first fruits development and every other week	2-3
Strawberry	3 – 5	From 1st fruits pre-maturity and every other week	1.5-2
Orchards, pone and stone fruits	3 – 5	Flower through fruiting	2-3
Citrus	1	Foliar apply 14 days from harvest	4
Citrus HLB Year 1 Program	2	Sept-Early Oct. then Nov-Early Dec.	2 - 4
Citrus HLB Maintenance	4	Spring, Summer, Sept., Nov	2 - 4
Young Tree Citrus HLB Maintenance	4	Spring, Summer, Sept., Nov.	3
Sugarcane	1	Foliar apply 14 days from harvest	4
Small Grains	1 – 3	Anytime, esp. at boot stage with fungicides	2-3
Corn, Soybeans, Cotton, Peanuts, Tobacco	1-3	Anytime from 5th leaf stage to 14 days prior to harvest	2-3

COMPATIBILITY AND STORAGE:

FERTILEADER Elite is compatible with many chemicals. In all cases, test the compatibility prior to treatment. Do not store under 32°F (0°C).

VOLUME: 264 Gallons (1000 liters) - NET WEIGHT: 3220 lb (1460 kg)

WARNING: Some crops may be injured by application of Boron.

Timac Agro USA, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Timac Agro USA, Inc. In no case shall Timac Agro USA, Inc. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the Buyer. Timac Agro USA, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

Timac Agro USÁ, INC. - P.O. Box 888 - Route 724 & Í-176 Reading, PA 19607-0888 – USA

MANUFACTURED BY TIMAC AGRO USA -P.O. Box 888 - Route 724 & I-176 Reading, PA 19607 -0888 - USA



FERTILEADER® Elite



WARNING

CAS: 15842-29-2

Contains: Calcium nitrate trihydrate

H302: Harmful if swallowed. H315: Causes skin irritation. H319: Causes serious eye irritation. H335: May cause respiratory irritation. P280: Wear face shield, protective clothing, protective gloves. P312: Call a POISON CENTER or doctor/physician if you feel unwell. P332+P313: If skin irritation occurs: Get medical advice/attention. P501: Dispose of this material and its container to hazardous or special waste collection point, in accordance with local, regional, national and/or international regulation.

Timac Agro USA, INC. - P.O. Box 888 - Route 724 & I-176 Reading, PA 19607-0888 - USA