

FERTILEADER®

Seactiv®

Gold

264 Gal.

FERTILEADER™ Gold

GUARANTEED ANALYSIS

- Water-soluble Boron (B).....5.7 %
 - Water-soluble Molybdenum (Mo).....0.35%
- DERIVED FROM: Boric Acid, Monoethanolamine, Sodium molybdate

F 1550

NON PLANT FOOD INGREDIENTS:

- Plant and seaweed extract 11 %

**PLEASE READ CAREFULLY BEFORE USING - KEEP OUT OF REACH OF CHILDREN
SHAKE WELL BEFORE USE**

DIRECTIONS FOR USE:

Do not spray during periods of frost. In hot conditions it is better to spray at the beginning or at the end of the day. Add FERTILEADER Gold in a half clean sprayer and agitate while mixing. Keep agitating until the end of the application. Dilute into at least 15 gallons/Acre. It can also be used in ULV aerial application. Foliar fertilization is intended as a supplement to a regular fertilization program and may not by itself provide all the nutrients normally required by crops or other intended plants. Warning some crops may be injured by application of Boron.

Crops	Number of applications	Application stages	Application rate (pints/Acre)
Alfalfa	2 - 3	After cutting with 2" to 6" of regrowth	2.5
Cole crops	1	Head formation	2.5
Sweet corn	2 - 3	At tassel	2
Peanuts	1	On third spray or at full bloom	2 - 3
Cotton	1	At pre-bloom to post-bloom	1 - 2
Peaches	1	At pre-bloom to post-bloom	1 - 2
Pecans	1	At pre-bloom to post-bloom	1 - 2
Onions	1 - 2	As needed for crop	2 - 3
Tobacco	1	As needed for crop	2 - 3
Potatoes	1	At mid to later crop stage	2 - 3
Soybeans	1	At pre-bloom to post-bloom	2 - 3
Grains, Grass seed crops	1	At flag leaf or head emergence	1 - 2
Cabbage, Cauliflower, Broccoli	1 - 3	From 10 days after transplantation until heads start developing	2 - 3
Green beans	1 - 2	From 3rd leaf to pre-bloom	1.5 - 3
Cucurbits	1	Apply at pre-bloom to post-bloom	1 - 3

COMPATIBILITY AND STORAGE:

FERTILEADER Gold is compatible with many chemicals. In all cases, test the compatibility prior to treatment. Do not store under 32°F (0°C).

Volume: 264 Gallons (1000 liters) - Net weight: 2733 lb (1230 kg)

TIMAC AGRO USA, INC. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of TIMAC AGRO USA, INC. In no case shall TIMAC AGRO USA, INC. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the Buyer. TIMAC AGRO USA, INC. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

MANUFACTURED BY TIMAC AGRO USA -
P.O. Box 888 - Route 724 & I-176 Reading,
PA 19607-0888 - USA





DANGER

CAS: 10043-35-3 / EC: 233-139-2

Contains: boric acid

H360: May damage fertility. P201: Obtain special instructions before use. P202: Do not handle until all safety precautions have been read and understood. P280: Wear protective clothing, face protection, protective gloves. P281: Use personal protective equipment as required. P308+P313: IF exposed or concerned: Get medical advice/attention. P501: Dispose of this material and its container to hazardous or special waste collection point, in accordance with local, regional, national and/or international regulation.