



# KSC-VI Phytactyl

## 14-12-14 Water Soluble Fertilizer

F 1550

**GUARANTEED ANALYSIS:**

Total Nitrogen (N).....	14.0%	Iron (Fe) water soluble.....	0.10%
13.0% Ammoniacal Nitrogen (N)		0.10% Iron (Fe)	
1.0% Nitrate Nitrogen (N)		Manganese (Mn) water soluble .....	0.1%
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) .....	12.0%	0.1% chelated Manganese (Mn)	
Soluble Potash (K <sub>2</sub> O) .....	14.0%	Zinc (Zn) water soluble .....	0.1%
Sulfur (S) .....	14.0%	0.1% chelated Zinc (Zn)	
Boron (B) water soluble .....	0.1%	Chlorine not more than .....	0.5%
Copper (Cu) water soluble .....	0.1%		
0.1% chelated Copper (Cu)			

DERIVED FROM: Ammonium Sulfate, Potassium Nitrate, Potassium phosphate, Potassium Carbonate, Borax Pentahydrate, Copper EDTA chelate, Iron EDTA chelate, Manganese EDTA Chelate, Zinc EDTA chelate.

NON PLANT FOOD INGREDIENTS: Seaweed extracts 0,7%.

**USE RECOMMENDATIONS:**

Crops	Number of Applications	Number of applications per Week	Application Stage	Rate
Golf Courses, General Turf and Sod	As Needed	As Needed	All Stages, Must immediately irrigate in	32 lbs/a provides .10 N/1000 78 lbs/a provides .25 lb N/1000 156 lbs/a provides .50 N/1000
Vegetables	2 to 15	1 to 5	From Transplant to post bloom	5 to 10 lbs per week
Row Crops, in furrow, in Liquid Solution	1	1	Applied in furrow at planting in solution, start	7.5 - 10 lbs/acre
Citrus, Nuts, Grapes	1	2 to 4	Use to start to supply Phosphorus quickly	25 lbs/500 gals water
Citrus, Nuts, Grapes Transplant Solution	2 to 10	1	At Transplant and 1-2 months	25 lbs/500 gals water

These recommendations are guidelines only. They should be adapted according to specific crop conditions, soil type and weather conditions. DILUTION: Dilute KSC VI Phytactyl in the mother solution with a concentration of 10-20% (22-44 lbs/26.4 gallons). This dilution should be considered as a guideline and have to be adapted in relation with the chemical characteristics of the irrigation water. Do not acidify mother solution below pH 6.0. COMPATIBILITY: KSC VI Phytactyl is compatible with most soluble fertilizers. Do not mix with products containing Calcium. Perform a prior test in all cases. Our responsibility is limited to guaranteeing the product in its original packaging. STORAGE: Store in a cool and dry place, away from heat and frost. Keep bag shut while not used. WARNING: Some crops may be injured by application of Boron.

**NET WEIGHT: 55.125 lbs. (25 kg)**

Timac Agro USA, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on the label when used in accordance with directions under normal use conditions. Crop Injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Timac Agro USA, Inc. In no case shall Timac Agro USA, Inc. be liable for consequential, special or indirect damages resulting from the use, handling, or shipping of this product. All such risk shall be assumed by the Buyer. Timac Agro USA, Inc. makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

Timac Agro USA, INC. - P.O. Box 888 - Route 724 & I-176 Reading, PA 19607-0888 - USA

DANGER - H360 - May damage fertility or the unborn child (oral). P201 - Obtain special instructions before use. P202 - Do not handle until all safety precautions have been read and understood. P280 - Wear protective clothing, eye protection, face protection. P308+P313 - If exposed or concerned: Get medical advice/attention. P405 - Store locked up. P501 - Dispose of contents/container to hazardous or special waste collection point, in accordance with local, regional, national and/or international regulation.

Manufactured by: Timac Agro España S.A. - Polígono Arazuri-Orcoyen - Calle C, nº32 - ORCOYEN (NAVARRA), 31160 ESPAÑA.

2EAKSCV/L-USA25 2239 - Made in Spain by Timac Agro España