



7-0-0

**TiMn IS A SPECIALTY LOW pH LIQUID NITROGEN, SULFUR AND MANGANESE
DESIGNED TO IMPROVE DEFICIENCIES AND CONDITION WATER.**

F1550

GUARANTEED ANALYSIS:

TOTAL NITROGEN (N)..... 7%
7% UREA NITROGEN

SULFUR (S)..... 5%

MANGANESE (Mn)..... 5%
5% WATER SOLUBLE MANGANESE

DERIVED FROM: UREA SULFATE, MANGANESE SULFATE.
NON PLANT FOOD INGREDIENTS: 2% SEAWEED EXTRACT,
98% OTHER INGREDIENTS

DIRECTIONS FOR USE: DO NOT SPRAY DURING PERIODS OF FROST. IN HOT CONDITIONS IT IS BETTER TO SPRAY AT THE BEGINNING OR AT THE END OF THE DAY. SHAKE WELL BEFORE USE.

FIELD, ROW, VEGETABLE, ORCHARD AND VINE CROPS: APPLY 1-4 QUARTS PER ACRE PER APPLICATION THROUGHOUT THE GROWING SEASON IN 15 TO 25 GALLONS OF WATER PER ACRE AS A DIRECTED FOLIAR SPRAY WHEN PLANTS ARE 8 TO 12 INCHES IN HEIGHT. TiMn CAN BE USED IN CONJUNCTION WITH GLYPHOSATE. ON AERIAL APPLICATIONS, USE MINIMUM 5 GAL/AC OF H₂O.

**KEEP OUT OF REACH OF CHILDREN - CAUTION -
READ ENTIRE LABEL BEFORE USING THIS PRODUCT**

CROPS	APPLICATION STAGE	APPLICATION RATE 1-6 APPLICATIONS
FIELD AND ROW CROPS, INCLUDING BUT NOT LIMITED TO: CORN, WHEAT, BARLEY, RYE, OATS, SORGHUM, RICE, OTHER CEREAL GRAINS, SUNFLOWER, CANOLA, SOYBEAN, LENTILS, DRY BEANS, FIELD PEAS, PEANUTS, COTTON, SUGARCANE	FOLIAR OR FERTIGATION APPLIED V3-V5 – EARLY VEG. (MONITOR SPRAY SOLUTION OR FERTILIZATION pH)	1 QT/AC FOLIAR 1-4 QT/AC SOIL APPLIED
ORCHARDS AND VINEYARDS INCLUDING BUT NOT LIMITED TO: APPLES, PEARS, PEACHES, NECTARINES, GRAPES (JUICE, TABLE WINE), CITRUS, BLUEBERRIES, HOPS	BUDBREAK TO PETAL FALL AND POST-HARVEST WITH ROW CLEANING PRODUCTS OR THROUGH FERTIGATION (MONITOR SPRAY SOLUTION OR FERTILIZATION pH)	1 QT/AC FOLIAR 1-4 QT/AC SOIL APPLIED
ROOT AND VEGETABLES, INCLUDING BUT NOT LIMITED TO: SUGAR BEETS, POTATOES, CARROTS, ONIONS, AND OTHER ROOT CROPS, TOMATOES, PEPPERS, EGGPLANTS, CUCURBITS, PUMPKINS, SQUASHES, CARROTS, STRAWBERRIES, CRANBERRIES, LEAFY GREENS, COLE CROPS (CABBAGE, BRUSSEL SPROUTS, BROCCOLI)	FOLIAR OR FERTIGATION APPLIED (MONITOR SPRAY SOLUTION OR FERTILIZATION pH)	1 QT/AC FOLIAR 1-4 QT/AC SOIL APPLIED
ALFALFA, CLOVER, GRASS CROPS, TURF	FOLIAR APPLY AT SPRING GREEN UP AND AGAIN 3-7 DAYS AFTER EACH CUTTING (MONITOR SPRAY SOLUTION OR FERTILIZATION pH)	1 QT/AC FOLIAR

THESE RECOMMENDATIONS ARE GUIDELINES ONLY. OPTIMUM RATE OF APPLICATION WILL VARY DEPENDING ON SOIL PROPERTIES (SUCH AS pH, ORGANIC MATTER CONTENT, TEXTURE), WEATHER CONDITIONS, TIME OF YEAR, PLANT SPECIES AND GENERAL HEALTH. FOR BEST RESULTS, FOLLOW SOIL AND TISSUE TEST RECOMMENDATIONS.

COMPATIBILITY, TANK MIXING AND STORAGE: CHECK MIXTURE COMPATIBILITY USING A JAR TEST PRIOR TO APPLICATION. FILL MIX TANK WITH HALF OF REQUIRED WATER AND BEGIN AGITATION. ADD PRODUCTS IN THIS ORDER, MIXING THOROUGHLY AFTER EACH ADDITION: TiMn, ADJUVANTS, PESTICIDES, THEN OTHER FERTILIZERS. FILL TANK WITH REMAINDER OF WATER AND CONTINUE AGITATION UNTIL SOLUTION IS COMPLETELY MIXED. MANUFACTURER RESPONSIBILITY IS LIMITED TO THE GUARANTEE OF THE PRODUCT'S COMPOSITION AS PER LABEL IN CERTIFIED PACKAGING. IN ALL CASES, TEST THE COMPATIBILITY PRIOR TO TREATMENT. STORE IN A COOL AND DRY PLACE, AVOID STORING IN SUNLIGHT OR HIGH HEAT ENVIRONMENTS. DO NOT STORE UNDER 32°F (0°C).

PRECAUTIONARY STATEMENTS: THIS IS A LOW pH PRODUCT. AVOID PROLONGED OR REPEATED CONTACT WITH EYES, SKIN, AND CLOTHING. CHEMICAL GOGGLES OR A FULL FACE SHIELD SHOULD BE WORN. TO PROTECT SKIN WEAR APPROPRIATE PROTECTIVE EQUIPMENT, SUCH AS RUBBER OR PLASTIC APRONS, RUBBER GLOVES, AND BOOTS. AVOID BREATHING MIST OR VAPOR. KEEP CONTAINERS CLOSED. WASH THOROUGHLY AFTER HANDLING. MAY CAUSE GASTROINTESTINAL DISTRESS IF SWALLOWED.

NET WEIGHT: 10.43 LB (4.73KG) PER GAL / 250 GAL. 2608 LB (1183KG)

TIMAC AGRO USA, INC. WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR PURPOSES STATED ON THE LABEL WHEN USED IN ACCORDANCE WITH DIRECTIONS UNDER NORMAL USE CONDITIONS. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF TIMAC AGRO USA, INC. IN NO CASE SHALL TIMAC AGRO USA, INC. BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE, HANDLING, OR SHIPPING OF THIS PRODUCT. ALL SUCH RISK SHALL BE ASSUMED BY THE BUYER. TIMAC AGRO USA, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

MANUFACTURED BY TIMAC AGRO USA
P.O. Box 888 - ROUTE 724 & I-176 READING, PA
19607 -0888 - USA



250 GAL