# Foliar Spray at Fourth Node

# Soybean

## OBJECTIVE

To assess the yield response of a foliar spray of Fertileader Gold at fourth node (V4) on crop of soybean.

Site Location:

Germansville, PA


Researcher: CMS, Inc. (Contract Research Org.)

#### **STUDY INFORMATION**

Variety Population S120114 Seedway 130,000 TIMAC AGRO PRODUCT



Soybean Yield Response from Fertileader Elite at Fourth Node (Germansville, PA)



### **KEY FINDINGS**

+3.9 bu/ac More than untreated control

> ROI: \$16.43/ac

**Graph:** Foliar spray of Fertileader Elite improved yield 3.9 bu/ac for soybean crop. The Gross Revenue above was calculated at \$8.50 bushel with Fertileader Elite retail cost of \$53.50/gallon.

## APPLICATION

| Treatment         | Application Rate |  |  |
|-------------------|------------------|--|--|
| Control           | N/A              |  |  |
| Fertileader Elite | 2.5 pint/A       |  |  |



#### **MATERIALS AND METHODS**

The experiment was implemented during the 2018 growing season at CMS, Inc., a private contract research organization. This location of the trial has been maintained weed and disease free, is well-drained, with excellent fertility. It has 1-2% slope but is well-suited to provide evenly distributed soil fertility, pH, soil organic matter, and water availability. Soil type is a rocky shale loam (Trexler shaly loam) with a pH of 6.8 (2.2% OM). Experimental units were plots four rows wide and 30 feet in length with 30-inch row spacing. Soybean was the previous crop and conventional tillage was used. A glyphosate tolerant variety (S120114 Seedway, Untreated) was grown at a population of 130,000 plants per acre to assess the impact of bio-nutritional products in an intensive crop system. Plots were arranged using an RCBD with 4 replications. Fertilizer applications were designed to be equivalent among all treatments and were applied pre-plant as a broadcast followed by incorporation. Fertileader Elite was applied at a rate of 2.5 pints/ac at fourth node (V4) growth stage. The center two rows of each plot were mechanically harvested at maturity for determination of grain yield and harvest moisture, and the yield was subsequently standardized to bushels/ac at 13% moisture.

#### **RESULTS AND CONCLUSIONS**

Foliar spray of Fertileader Elite (2.5 pint/A) at fourth node improved soybean yield over untreated control by 3.9 bushel/acre. This resulted in a ROI of \$16.43/acre.

| Treatment                       | Yield<br>bu/ac | Gross Revenue @<br>\$8.50/bu | Change<br>from<br>Control | Added<br>Costs/ac | ROI     |
|---------------------------------|----------------|------------------------------|---------------------------|-------------------|---------|
| Control                         | 40.8           | \$346.80                     | -                         | \$0.00            | -       |
| Fertileader Elite<br>(2.5 pt/A) | 44.7           | \$379.95                     | \$33.15                   | \$16.72           | \$16.43 |

### **RETURN ON INVESTMENT**

**İmac** Agro | **R&D** 

Author: Alex Duffy, National Product Manager aduffy@timacusa.com 484-869-3043 (please contact if further information is needed)

3/17/2021

us.timacagro.com | 800-545-5474